



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Beyond land cover change: Towards a new generation of Land Use Models

Citation for published version:

Verburg, PH, Alexander, P, Evans, T, Magliocca, NR, Malek, Z, Rounsevell, M & van Vliet, J 2019, 'Beyond land cover change: Towards a new generation of Land Use Models', *Current Opinion in Environmental Sustainability*, vol. 38, pp. 77-85. <https://doi.org/10.1016/j.cosust.2019.05.002>

Digital Object Identifier (DOI):

[10.1016/j.cosust.2019.05.002](https://doi.org/10.1016/j.cosust.2019.05.002)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Current Opinion in Environmental Sustainability

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Beyond land cover change: Towards a new generation of Land System Models

Peter H. Verburg^{1,2*}, Peter Alexander^{3,4}, Tom Evans⁵, Nicholas R. Magliocca⁶, Ziga Malek², Mark D.A. Rounsevell^{3,7}, Jasper van Vliet²

¹Federal Institute for Forest, Snow and Landscape Research (WSL), Zürcherstrasse 111, CH-8903 Birmensdorf, Switzerland

²Institute for Environmental Studies, Vrije Universiteit Amsterdam, de Boelelaan 1087, 1081HV Amsterdam, the Netherlands

³School of GeoSciences, University of Edinburgh, Drummond Street, Edinburgh, UK

⁴Global Academy of Agriculture and Food Security, The Royal (Dick) School of Veterinary Studies, University of Edinburgh, Midlothian, UK

⁵School of Geography and Development, University of Arizona, United States of America

⁶Department of Geography, University of Alabama, Tuscaloosa, Alabama, United States of America

⁷Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research, Atmospheric Environmental Research (IMK- IFU), Garmisch-Partenkirchen, Germany

*Corresponding author, peter.verburg@vu.nl

Highlights:

- Lack of new land use or land system modelling concepts published in recent year
- Land system models are frequently insufficiently evaluated
- Advances in better representation of human agency in land system models are identified
- Large potential identified for land system models to contribute to the identification and design of sustainability solutions

28

29 *Abstract*

30 Land system models play an important role in exploring future land change dynamics and are
31 instrumental to support the integration of knowledge in land system science. However, unsatisfactory
32 progress has been made in achieving these aims due to insufficient model evaluation and limited
33 representation of the underlying socio-ecological processes and human agency. We discuss how land
34 system models can better represent multi-scalar dynamics, human agency and demand-supply
35 relations, as well as how to achieve deeper insights from model evaluation. By addressing these issues
36 we outline pathways towards a new generation of land system models that allow not only the
37 assessment of future land cover pattern changes, but also enable societal envisioning in supporting
38 the design of sustainability solutions.

39

40 *Keywords:* Land use, Land cover, Simulations, Integrated Assessment, Agent-based models,
41 Sustainability

42

43 **Introduction**

44 As with other emerging scientific fields, rapid advances in land use modelling were made during the
45 first decades of the development of land system science. Several alternative paradigms for modelling
46 land use change processes were developed [1]. Over the past decade, the number of publications
47 related to land use change modelling has continued to increase. Publications in this period indicate
48 three trends: 1) The frequent application of easily available land use models in case-studies aimed at
49 informing spatial planning. Many of these studies apply relatively simple spatial models, e.g. using a
50 combination of Markov chains for the quantity of change and cellular automata to emulate patterns
51 of land cover change [2]; 2) The incremental improvement of existing models and modelling concepts
52 [3-5]; 3) The development of agent-based models for specific case-studies that are difficult to
53 generalize beyond the specific context, characterized by O'Sullivan et al. [6] as the YAAWN syndrome
54 ("Yet Another Agent-Based Model ... Whatever ... Nevermind ...").

55 These trends illustrate the relevance of land change modelling as a tool in land system science.
56 However, the limited amount of novel modelling concepts put forward raises the question of whether
57 current tools and modelling concepts allows the full potential of land system science to be reached?

58 This question cannot be answered in a generic manner. In many projects existing land use models
59 have successfully played a role in synthesizing project results [7,8] or in structuring discussions with
60 stakeholders [9,10]. In spite of this, there is a recurring notion in the literature of the model being
61 presented as the endpoint, rather than as a process of learning from the model design or application.
62 This is unfortunate since modelling systems are rarely adopted by stakeholders after the lifetime of a
63 project [11].

64 The application of currently available models for policy and planning is hampered by uncertainty
65 throughout the modelling process, and limited progress has been reported in understanding or
66 reducing this uncertainty. While predictive accuracy is just one metric of a model's value, earlier
67 validation efforts showed that few land use models outperformed a simple 'no change' model [12].
68 More recently, Mas et al. [13] showed that for a similar (virtual) landscape four different, frequently
69 used, land change models resulted in strongly different outcomes. A comparison of global land use

models and integrated assessment models (IAMs) showed that, in essence, the differences between the models analyzed were greater than the differences between the different scenarios modelled [14,15]. A review of calibration and validation practices in land use models [16] found that 31% of the applications did not report any model evaluation, while the rest were predominantly assessed in terms of their location accuracy, ignoring the uncertainty in the quantity and spatial patterns of land use. Only 17% of the model applications reported an uncertainty analysis, and 12% reported a sensitivity analysis.

Given these conditions, the objective of this paper is to identify opportunities to improve land use modelling towards a new generation of land system models that is better able to synthesize and formalize insights, make sources of uncertainty in projections transparent, and support the design of sustainability solutions.

Key dimensions for land system modelling

Addressing the multi-scalar challenge

The dilemma of choosing an appropriate scale for modelling is well-known for land system science and multi-scalar dynamics have been a challenge since the origin of the research field [20-22]. Global drivers affect places in different ways and aggregate impacts of local responses feedback to the global system. Coupling of models operating at different scales has been proposed to address the multiple levels of analysis needed to describe all important processes [23-26]. However, usually only a one-way, top-down flow of data occurs, as incorporating feedbacks would lead to computationally complex iterations between models. Moreover, different modelling concepts and behavioral assumptions at different levels may lead to inconsistencies between models at different scales (“ugly constructs” according to Voinov and Shugart [27]).

Addressing the multi-scalar challenge requires new model structures that are truly multi-scale, and thus likely more complex than current models. Experiences in other fields of science may be instrumental to inform such a design, i.e. the multi-level structure employed in remote sensing [27a] or multi-scale modelling in physics [27b]. In physics sometimes a sequential modelling is used in which micro-models pre-compute details of some of the constitutive relations in the macro-model. Such an approach may also be used in land system science. In addition, rather than simulating all underlying processes, a larger role may be given to meta-studies, synthesizing empirically measured responses in local studies to inform model design [28]. Such micro-level models or meta-studies should aim to synthesize the role of contextual conditions on responses, which can be translated into simple model rules implemented within a higher level model to account for the local level responses.

When feedbacks between micro and macro levels are important, concurrent multi-scale modelling (also referred to as nested modelling) may be applied in which quantities needed in the macro-scale model are computed on-the-fly from micro-scale models as the computation proceeds. Concurrent coupling allows one to evaluate these forces at the locations where they are needed to resolve local behavior and then use macro-models elsewhere.

Conversely, new models could focus on different model structures that better reflect the important scalar dynamics more explicitly. For example, models of global trade flows of food commodities use a different approach than models of decision making about land use at local levels. However, combining models of trade flows with sub-national models of human agency would create new modelling approaches of considerable utility. An example of such a modelling approach is provided by Lamperti

et al. [30] who provide an alternative, agent-based, model structure to the classic coupling of general equilibrium models and climate models in IAMs.

A confounding factor in the multi-scalar challenge is the wide variety of telecoupled processes and the impact of location conditions on land change outcomes [31,32]. The location of agricultural expansion will affect production and impacts on biodiversity. Extent and spatial patterns of expansion will differ between locations as a result of the local socio-economic, cultural and demographic context. While downscaling of global model outcomes to pixels is well established and an integral part of IAMs, feedbacks from the local to the global-level are poorly captured with this approach. Global land use models are, therefore, often unable to appropriately capture processes such as displacement effects, multi-level governance of land use, adaptive learning, not fully economically driven decision making and human behavior that underpins decision making [32]. Part of such bottom-up processes could be captured by nesting micro-models within the macro-models to capture bottom-up responses in a more adequate manner as has been described above. However, when these responses are moderated through processes at different scales capturing the bottom-up response may not be sufficient. Displacement and other spill-over effects can occur through multiple mechanisms [55][32a] and the spatial scale across which these effects occur depends on the actors and processes involved, such as the structure of the value chain and markets, which can cause spillovers to occur from within the same landscape to across world regions. Economic models (i.e. equilibrium models) that can address such displacement processes have fixed representations of modelling units and can only address displacement effects between these units. While qualitative methods and conceptual models are able to describe such cross-scale mechanisms, the consistent representation in models is still challenging.

Resilience of the land system has received increased attention given the interconnectedness of the global system and the potential consequences from shocks, such as extreme weather events or trade conflicts. Land system models have potential to simulate such shocks and the resulting consequence so that the systems behaviors can be better understood and negative outcomes mitigated. However, to date such modelling remains lacking, in part due to the challenges in consider cross scale interaction in dynamic (i.e. non-equilibrium) conditions.

The next generation of land use change models should give more attention to cross-scale processes rather than focusing on a single scale or level of modelling to better represent spillover processes that are increasingly important in globalized land use. To achieve this we may have to move away from using simple spatial (i.e. world regions or pixels) or organizational entities (individual or institutional agents) as units of simulation to more blended approaches where the processes of interaction are central to the simulation rather than the units affected by these processes.

Embrace complexity and diversity of human agency

The attractiveness of agent-based modelling for land use change originates from the explicit representation of the diversity in decision making and the desire to incorporate agent-interactions [33,34]. However, most agent-based modelling is focused on the local scale because finding sufficient empirical data about decision making processes and outcomes at larger scales is extremely difficult. Attempts to use agent-based models at larger scales often resort to simplifications of the variation in decision making by linking agent-types directly to land cover types. While there is general agreement that decision making dynamics in land use can vary strongly across the globe, there is little empirical basis or theoretical insight to help selecting from different approaches to represent decision-making in simulation models [35].

To fill this gap, Malek et al. (subm) conducted a meta-analysis of case-studies to identify where, and under what conditions, certain modes of land-use decision making are found. The occurrence of archetypical types of decision making, ranging from satisficing behavior to utility maximization, were related to contextual conditions, leading to a predictive model that indicates what mode of land use decision making can be expected in a particular context (Figure 1). In spite of the large generalization, this synthesis is a first step towards global land use models that represent the variation in decision making [32]. In addition to the spatial variations, land-use decision making often shows an evolution over time [36], as is represented in models that incorporate adaptive behavior [37]. Moving away from the assumption of uniform and static decision making is a big step for large-scale land use models and does not necessarily mean that all should become agent-based modelling and represent individual agents. Differences in decision making mechanisms can also be reflected in spatial models that use pixels as units of simulation, either through the choice of determinants of location suitabilities or through the spatial extent accounted for in choosing the most optimal location for a particular land use. In large-scale global economic and integrated assessment models there have been several calls to represent heterogeneity and some early approaches have been proposed, however not yet related to land use [37a][37b].

A limitation of most agent-based modelling in land use studies is the focus on primary actors of land use change, mostly farmers. Recent developments show an increasing influence on land use decisions of distant land owners, investors and companies through large-scale land acquisitions, contract farming and investments [38,39]. To better account for such developments, insight into the decision making of these actors needs to be obtained. There are yet few studies [39a][39b] that explicitly account for this type of land change dynamics as most existing models are more tailored towards representing small-holder farmer decisions.




Figure 1: Results of a meta-analysis of case studies reporting decision-making modes worldwide. Left: Radar charts showing average scores on abilities (financial, land size, land tenure, connectedness, power), objectives (survival, economic, environmental, lifestyle, social prestige) and attitudes (change, legislation, environmental values) for the different decision-modes; Right: Maps depicting likelihood of finding a specific decision mode based on extrapolation with socio-economic and biophysical context variables (Malek et al. subm).

Linking demand and supply

While land system science is based on the notion of socio-ecological systems, we often still model only a symptom of the socio-ecological system dynamics: the conversion of one land cover to another. This is a direct result of the dependence on remote sensing data that reflect land cover. Only few models account for the, historically, most important pathway of fulfilling increasing demands for land-based commodities: land use intensification [36,40,41]. Instead, most models assume an external pressure

or demand to steer land cover change quantities or use Markov chains to extrapolate from historic trends. The processes underlying this demand are not modelled explicitly and feedbacks between demand and supply are ignored. As an exception, general economic equilibrium models determine demand and supply across the full economy, where the costs of production can affect consumption patterns through price signals [42] and an economic choice is made between expansion of land area or intensification. Such analysis is useful, but lacks a representation of the spatial heterogeneity, as the spatial resolution of these models is often restricted to world regions. Additionally, other feedbacks in the system, including lifestyles, land tenure, advertising, markets and governance, can only be incorporated in stylized forms, e.g., through demand elasticities and production costs. Consumption of agricultural commodities, but also the use of other land-based commodities such as biofuels, are strongly determined by large corporations that impact consumption choices through markets and price incentives. Governments impact relationships between demand and supply, e.g., through trade barriers and subsidies for production or export of products [43]. As a result, consumer prices do not reflect the real production costs and consumer choices are often not economically rational, let alone fully accounting for health or environmental costs and benefits. While it is generally acknowledged that consumption choices are a strong determinant of land system change and offer a large potential to reduce pressures on land resources and environmental impact [44-46], there is relatively limited attention to this in land use modelling. The links and feedbacks between consumer and producer choices need to be represented in modelling to avoid a sole focus on production side solutions. Transformative change towards sustainability requires also addressing the fundamental drivers and behavioral choices in socio-ecological systems that are the underlying causes of land system changes. While there is a rich literature on consumption behavior and an emerging knowledge on the role of supply chains [46a] these are hardly captured in land system models to allow a more quantitative exploration of the different pathways to fulfilling the demand for land-based products more sustainably.

Learning from modelling

Land use modelling is too often presented as a goal in itself, or for the purpose of 'prediction'. However, the model building and testing process is most useful in advancing our understanding of land use systems. This is evident in the rapidly growing field of participatory modelling, which aims at engaging the knowledge of stakeholders to create formalized and shared representations of reality and using models as boundary objects to collectively reason about environmental problems and foster two-way learning [Jordan et al. 2018][Gray et al 2018]. A review of 180 environmental sciences papers using participatory modelling identified a gap between the qualitative and quantitative development phases that hampers the use of participatory approaches to develop the more quantitative models for scenario analysis (Voinov et al. ??). However, final results are typically not the most valuable or convincing aspects of a modelling effort for policy makers and other stakeholders, but rather the rationale of the (cascading) processes of impact of a certain intervention [9,47]. Rather than seeing the model as a black box, it is the internal logic leading to a specific outcome that needs to be uncovered to convince stakeholders of appropriate actions. Similarly, models are also an important learning tool for researchers. Models force us to formalize our understanding of land systems: select those processes that are important, quantify relations and bring different components together into a consistent whole. In that sense, models can be a boundary object (i.e., platform that spans disciplinary boundaries to enable contributions and interpretations from diverse perspectives) in socio-ecological systems analysis [48].

Comparing model outcomes against reality (i.e., model validation) is also an opportunity for improving our system understanding [49]. While model validation is rather common for local to regional scale models [16], most global land use models have still never been compared against data [50]. Until recently, global land cover products were of insufficient quality to enable full validation. Recent global, multi-temporal datasets offer new opportunities to validate global land change models [51]. Of course, validation based on land cover outcomes is not necessarily conclusive, because different land change processes may lead to the same patterns (equifinality) and calibration based on past conditions does not imply predictability of future conditions. However, model evaluation, which includes validation, as well as uncertainty analysis, model verification, sensitivity analysis, and benchmarking (comparison with other models), is an essential step in learning about the system and the range of applications the model is suited for.

Recent model comparisons [14,15] show that large differences in outputs exist between land use models, even though most of the compared models use a common modelling paradigm (*viz.* IAMs). As these models are used to inform large-scale governmental assessments, such as those of the IPCC and to a lesser extent IPBES, this uncertainty is concerning. In addition, land use results in these assessments are harmonized from only one of many possible land-use models [52], and then used by climate or ecosystem models to explore uncertainty [53] - an approach which may neglect key elements of uncertainty in the land use projections. Furthermore, separate scenarios have been assigned to different individual models or a small group of IAMs [54], carrying the risk that urgent policy decisions are based on information that hardly reflects the uncertainty embedded in the choice of model.

While large differences between different model types are a challenge from a predictive point of view, they provide an opportunity for learning from model comparisons. Models that can simultaneously implement multiple, alternative process representations provide a computational laboratory to explore the applicability of hypothesized land system processes across a range of conditions, and iteratively improve our understanding of the broader socio-ecological system. Model representations that balance specificity and generality are a tool for theory development and testing, particularly for middle-range theories [55]. This approach is exemplified by Magliocca et al. [56] who tested the validity of generic theory to explain land use changes across different contexts in a virtual laboratory setting.

To derive the greatest insights from models, results need to be repeatable by researchers outside of the groups where a model was developed. This requires comprehensive model descriptions and full scenario outputs to be published, as well as making model code available with complete sets of input data to allow re-running or adaptation of existing simulations

Moving beyond exploration: land use modelling for the envisioning and design of sustainable futures

The majority of land change models are used to project exploratory scenarios under assumed future conditions. While such scenario studies have proven useful in anticipating future land use outcomes under uncertain drivers, it is often difficult to link these to the policy, behavioral and management decisions needed to arrive at more beneficial outcomes. As most of the model structures are based on current processes and parameterized or calibrated on past or current conditions, these models are not suited to assess socio-ecological system developments that strongly deviate from past conditions, such as the impacts of de-growth [57] or large scale migration [58] on land use. At the same time, awareness is growing that meeting the sustainable development goals requires large societal

transformations, including behavioral changes, technological shifts and institutional arrangements. Most models are only able to address the ‘shallow leverage points’ of sustainability transformations and lack the capacity to address ‘deep leverage points’ [58a]. Moreover, such sustainability transformations will come with significant tradeoffs that require far-reaching decisions and societal envisioning processes.

Land use models have the capability to support societal envisioning processes by sketching out the land use realities of alternative objectives and quantifying the tradeoffs associated with those [59]. Modelling can help to explore land use futures that navigate such tradeoffs by optimizing sets of objectives while minimizing tradeoffs [46,60,61]. Examples that move beyond exploratory scenario modelling include the work of Wolff et al. [62] that visualized how the world would look like if all agreed land restoration goals in international treaties were met, and Mehrabi et al. [63] who assess the consequences of conserving half of the land area for biodiversity conservation. While the individual goals of these studies are laudable, the resulting global land use patterns may not be considered the most desirable due to competing claims for space. Such studies help the translation of single goals to more consistent and synergetic land use futures and open the debate on what future land use we want. Verkerk et al. [64] sketch an alternative approach where stakeholder visions are matched with a large set of exploratory scenarios to identify the conditions and policies that would bring land use closer to stakeholder defined visions. This paper is an early implementation of the call of Rosa et al. [65] for visionary biodiversity scenarios. A more advanced implementation of this approach is presented by Cooper and Dearing [66] who model fishery systems to show under which conditions different pathways to safe and just socio-ecological systems are feasible.

Conclusion

Land use modelling can play multiple roles within land system science and has a critical role in major environmental assessments, both as a mechanism to evaluate drivers of global environmental change and as a means to mitigate or adapt to global change. While progress is made in refining existing models and in the field of participatory modelling, major aspects important in representing socio-ecological system dynamics are insufficiently addressed in existing models and many approaches do not use comprehensive model evaluation procedures to secure an understanding of uncertainty and a continuous learning process. While these challenges are mentioned before, progress towards resolving these is small. Land system modelling needs to move beyond incremental improvements towards testing new model structures and new workflows focused on cross-scale interactions, diversity in human agency and the links between demand and supply sites. This may lead to increased complexity of models which conflicts with calls for simpler models to support stakeholder engagement and inform policy making. However, lower complexity does not mean better science, and simplifications can lead to potentially incorrect conclusions if spill-overs and feedbacks are ignored. We thus argue that while higher complexity models may be more difficult to use in policy circles, such difficulty is clearly offset by their greater realism and rigor [37b]. Currently, many important aspects of land system science are only addressed by qualitative methods and ignored by large scale models used at the science-policy interface (i.e. IPCC and IPBES). Therefore, we call upon the scientific community for innovative modelling approaches that better embed our understanding of land systems, and on the lead scientists of major assessments such as IPCC and IPBES to move beyond the established set of IAMs and open up to insights obtained from new land system model types. This way, land system science could move beyond using models as assessment tools and towards the use of models as virtual laboratories to stimulate societal learning and the co-design of sustainability solutions.

Acknowledgements

PV would like to acknowledge funding from European Research Council grant GLOLAND (Grant No. 311819). PA would like to acknowledge support by the UK's Global Food Security Programme project Resilience of the UK food system to Global Shocks (BB/N020707/1). NM would like to acknowledge the support of an award from the U.S. National Science Foundation EAGER ISN # #1837698. MR would like to acknowledge the support of the European Commission funded IMPRESSIONS project (FP7).

References

- 1.* Brown DG, Verburg PH, Pontius Jr RG, Lange MD: **Opportunities to improve impact, integration, and evaluation of land change models**. *Current Opinion in Environmental Sustainability* 2013, **5**:452-457.
Paper describing the different roles of land change models in different phases of a policy cycle.
2. Eastman JR: **IDRISI Kilimanjaro guide to GIS and image processing**. Edited by. Worcester MA: Clark Labs; 2003.
3. Gounaridis D, Chorianopoulos I, Symeonakis E, Koukoulas S: **A Random Forest-Cellular Automata modelling approach to explore future land use/cover change in Attica (Greece), under different socio-economic realities and scales**. *Science of The Total Environment* 2019, **646**:320-335.
4. Liang X, Liu X, Li D, Zhao H, Chen G: **Urban growth simulation by incorporating planning policies into a CA-based future land-use simulation model**. *International Journal of Geographical Information Science* 2018, **32**:2294-2316.
5. Yang X, Zheng XQ, Chen R: **A land use change model: Integrating landscape pattern indexes and Markov-CA**. *Ecological Modelling* 2014, **283**:1-7.
- 6.* O'Sullivan D, Evans T, Manson S, Metcalf S, Ligmann-Zielinska A, Bone C: **Strategic directions for agent-based modeling: avoiding the YAAWN syndrome**. *Journal of Land Use Science* 2016, **11**:177-187.
Critical reflection on progress in agent-based modelling pointing towards weaknesses in current practice
7. Harrison PA, Dunford R, Savin C, Rounsevell MDA, Holman IP, Kebede AS, Stuch B: **Cross-sectoral impacts of climate change and socio-economic change for multiple, European land- and water-based sectors**. *Climatic Change* 2015, **128**:279-292.
8. Lotze-Campen H, Verburg PH, Popp A, Lindner M, Verkerk PJ, Moiseyev A, Schrammeijer E, Helming J, Tabeau A, Schulp CJE, et al.: **A cross-scale impact assessment of European nature protection policies under contrasting future socio-economic pathways**. *Regional Environmental Change* 2018, **18**:751-762.
- 9.** Voinov A, Kolagani N, McCall MK, Glynn PD, Kragt ME, Ostermann FO, Pierce SA, Ramu P: **Modelling with stakeholders – Next generation**. *Environmental Modelling & Software* 2016, **77**:196-220.
Strong overview of the ways in which models can support stakeholder engagement.
10. Zagaria C, Schulp CJE, Kizos T, Gounaridis D, Verburg PH: **Cultural landscapes and behavioral transformations: An agent-based model for the simulation and discussion of alternative landscape futures in East Lesvos, Greece**. *Land Use Policy* 2017, **65**:26-44.
11. Zasada I, Piorr A, Novo P, Villanueva AJ, Valánszki I: **What do we know about decision support systems for landscape and environmental management? A review and expert survey within EU research projects**. *Environmental Modelling & Software* 2017, **98**:63-74.

12. Pontius R, Boersma W, Castella J-C, Clarke K, de Nijs T, Dietzel C, Duan Z, Fotsing E, Goldstein N, Kok K, et al.: **Comparing the input, output, and validation maps for several models of land change.** *The Annals of Regional Science* 2008, **42**:11-37.
13. Mas J-F, Kolb M, Paegelow M, Camacho Olmedo MT, Houet T: **Inductive pattern-based land use/cover change models: A comparison of four software packages.** *Environmental Modelling & Software* 2014, **51**:94-111.
14. ** Alexander P, Prestele R, Verburg PH, Arneth A, Baranzelli C, Batista e Silva F, Brown C, Butler A, Calvin K, Dendoncker N, et al.: **Assessing uncertainties in land cover projections.** *Global Change Biology* 2017, **23**:767-781.
Reports a comparison of land use outcomes of Global and European scale land use land use and integrated assessment models. Shows clearly the large uncertainties and differences between models.
15. ** Prestele R, Alexander P, Rounsevell MDA, Arneth A, Calvin K, Doelman J, Eitelberg DA, Engström K, Fujimori S, Hasegawa T, et al.: **Hotspots of uncertainty in land-use and land-cover change projections: a global-scale model comparison.** *Global Change Biology* 2016, **22**:3967-3983.
Analyzes uncertainties between Global scale land use models in a spatial manner and identifies critical regions of uncertainty.
16. * van Vliet J, Bregt AK, Brown DG, van Delden H, Heckbert S, Verburg PH: **A review of current calibration and validation practices in land-change modeling.** *Environmental Modelling and Software* 2016, **82**:174-182.
Review of current practice in land change model evaluation.
20. Evans T, Ostrom E, Gibson C: **Scaling Issues With Social Data in Integrated Assessment Modeling.** *Integrated Assessment* 2003, **3**:135-150.
21. Gibson CC, Ostrom E, Anh TK: **The concept of scale and the human dimensions of global change: a survey.** *Ecological Economics* 2000, **32**:239.
22. Veldkamp A, Verburg PH, Kok K, de Koning GHJ, Priess J, Bergsma AR: **The need for scale sensitive approaches in spatially explicit land use change modeling.** *Environmental Modeling and Assessment* 2001, **6**.
23. Millington J, Xiong H, Peterson S, Woods J: **Integrating Modelling Approaches for Understanding Telecoupling: Global Food Trade and Local Land Use.** *Land* 2017, **6**:56.
24. Verburg PH, Tabeau A, Hatna E: **Assessing spatial uncertainties of land allocation using a scenario approach and sensitivity analysis: A study for land use in Europe.** *Journal of Environmental Management* 2013, **127**:S132-S144.
25. Verstegen JA, Hilst F, Woltjer G, Karssenbergh D, Jong SM, Faaij APC: **What can and can't we say about indirect land-use change in Brazil using an integrated economic – land-use change model?** *GCB Bioenergy* 2016, **8**:561-578.
26. ** Robinson DT, Di Vittorio A, Alexander P, Arneth A, Barton CM, Brown DG, Kettner A, Lemmen C, O'Neill BC, Janssen M, et al.: **Modelling feedbacks between human and natural processes in the land system.** *Earth Syst. Dynam.* 2018, **9**:895-914.
Provides an overview of methods for coupling models with a focus on coupling human and natural processes illustrated with examples.
27. Voinov A, Shugart HH: **'Integronsters', integral and integrated modeling.** *Environmental Modelling & Software* 2013, **39**:149-158.
- 27a. Watmough GR, Marcinko CLJ, Sullivan C, Tschirhart K, Mutuo PK, Palm CA, Svenning J-C: **Socioecologically informed use of remote sensing data to predict rural household poverty.** *Proceedings of the National Academy of Sciences* 2019, **116**:1213.
- 27b. Weinan E: *Principles of multiscale modeling*: Cambridge University Press; 2011.
28. Magliocca NR, van Vliet J, Brown C, Evans TP, Houet T, Messerli P, Messina JP, Nicholas KA, Ornetsmüller C, Sagebiel J, et al.: **From meta-studies to modeling: Using synthesis knowledge to build broadly applicable process-based land change models.** *Environmental Modelling & Software* 2015, **72**:10-20.

- 30.** Lamperti F, Dosi G, Napoletano M, Roventini A, Sapio A: **Faraway, So Close: Coupled Climate and Economic Dynamics in an Agent-based Integrated Assessment Model.** *Ecological Economics* 2018, **150**:315-339.
Interesting example of a study presenting an alternative model structure and concept, leading to strongly deviating outcomes.
31. Friis C, Nielsen J: **On the System. Boundary Choices, Implications, and Solutions in Telecoupling Land Use Change Research.** *Sustainability* 2017, **9**:974.
32. Rounsevell MDA, Arneth A, Alexander P, Brown DG, de Noblet-Ducoudré N, Ellis E, Finnigan J, Galvin K, Grigg N, Harman I, et al.: **Towards decision-based global land use models for improved understanding of the Earth system.** *Earth Syst. Dynam.* 2014, **5**:117-137.
- 32a. Liu J, Dou Y, Batistella M, Challies E, Connor T, Friis C, Millington JDA, Parish E, Romulo CL, Silva RFB, et al.: **Spillover systems in a telecoupled Anthropocene: typology, methods, and governance for global sustainability.** *Current Opinion in Environmental Sustainability* 2018, **33**:58-69.
33. Manson SM: **Simplifying complexity: a review of complexity theory.** *Geoforum* 2001, **32**:405-414.
34. Murray-Rust D, Brown C, van Vliet J, Alam SJ, Robinson DT, Verburg PH, Rounsevell M: **Combining agent functional types, capitals and services to model land use dynamics.** *Environmental Modelling & Software* 2014, **59**:187-201.
- 35.* Müller-Hansen F, Schlüter M, Mäs M, Donges JF, Kolb JJ, Thonicke K, Heitzig J: **Towards representing human behavior and decision making in Earth system models – an overview of techniques and approaches.** *Earth Syst. Dynam.* 2017, **8**:977-1007.
Good overview paper of decision making mechanisms and their implementation in Earth system models.
36. Brown C, Holzhauser S, Metzger MJ, Paterson JS, Rounsevell M: **Land managers' behaviours modulate pathways to visions of future land systems.** *Regional Environmental Change* 2018, **18**:831-845.
37. Holman IP, Brown C, Carter TR, Harrison PA, Rounsevell M: **Improving the representation of adaptation in climate change impact models.** *Regional Environmental Change* 2018.
- 37a. McCollum DL, Wilson C, Pettifor H, Ramea K, Krey V, Riahi K, Bertram C, Lin Z, Edelenbosch OY, Fujisawa S: **Improving the behavioral realism of global integrated assessment models: An application to consumers' vehicle choices.** *Transportation Research Part D: Transport and Environment* 2017, **55**:322-342.
- 37b. Mercure J-F, Pollitt H, Bassi AM, Viñuales JE, Edwards NR: **Modelling complex systems of heterogeneous agents to better design sustainability transitions policy.** *Global Environmental Change* 2016, **37**:102-115.
38. Jayne TS, Chamberlin J, Headey DD: **Land pressures, the evolution of farming systems, and development strategies in Africa: A synthesis.** *Food Policy* 2014, **48**:1-17.
39. Otsuka K, Nakano Y, Takahashi K: **Contract Farming in Developed and Developing Countries.** *Annual Review of Resource Economics* 2016, **8**:353-376.
- 39a. Hailegiorgis AB, Cioffi-Revilla C: **Agent-Based Modeling of Large-Scale Land Acquisition and Rural Household Dynamics.** In *GeoComputational Analysis and Modeling of Regional Systems.* Edited by Thill J-C, Dragicevic S: Springer International Publishing; 2018:101-119.
- 39b. Debonne N, van Vliet J, Verburg PH: **Future governance options for large-scale land acquisition in Cambodia: Impacts on tree cover and tiger landscapes.** *Environmental Science & Policy* 2019, **94**:9-19.
40. Asselen S, Verburg PH: **Land cover change or land use intensification: simulating land system change with a global-scale land change model.** *Global Change Biology* 2013, **19**:3648-3667.
41. Alexander P, Rabin S, Anthoni P, Henry R, Pugh TAM, Rounsevell MDA, Arneth A: **Adaptation of global land use and management intensity to changes in climate and atmospheric carbon dioxide.** *Global Change Biology* 2018, **24**:2791-2809.

42. Lampe M, Willenbockel D, Ahammad H, Blanc E, Cai Y, Calvin K, Fujimori S, Hasegawa T, Havlik P, Heyhoe E, et al.: **Why do global long-term scenarios for agriculture differ? An overview of the AgMIP Global Economic Model Intercomparison.** *Agricultural Economics* 2014, **45**:3-20.
43. Anderson KV, Ernesto Martin, Will: *The Relative Importance Of Global Agricultural Subsidies And Market Access*, vol 5569. Available at SSRN: <https://ssrn.com/abstract=912473>; 2006.
44. Alexander P, Rounsevell MDA, Dislich C, Dodson JR, Engström K, Moran D: **Drivers for global agricultural land use change: The nexus of diet, population, yield and bioenergy.** *Global Environmental Change* 2015, **35**:138-147.
45. Springmann M, Wiebe K, Mason-D'Croz D, Sulser TB, Rayner M, Scarborough P: **Health and nutritional aspects of sustainable diet strategies and their association with environmental impacts: a global modelling analysis with country-level detail.** *The Lancet Planetary Health* 2018, **2**:e451-e461.
- 46.* Van Kernebeek HRJ, Oosting SJ, Van Ittersum MK, Bikker P, De Boer IJM: **Saving land to feed a growing population: consequences for consumption of crop and livestock products.** *The International Journal of Life Cycle Assessment* 2016, **21**:677-687.
An interesting approach using modelling to determine optimal diets for limiting land use competition.
- 46a. Lambin EF, Gibbs HK, Heilmayr R, Carlson KM, Fleck LC, Garrett RD, le Polain de Waroux Y, McDermott CL, McLaughlin D, Newton P, et al.: **The role of supply-chain initiatives in reducing deforestation.** *Nature Climate Change* 2018, **8**:109-116.
47. Ferré TPA: **Revisiting the Relationship Between Data, Models, and Decision-Making.** *Groundwater* 2017, **55**:604-614.
48. Reilly AC, Dillon RL, Guikema SD: **Agent-Based Models as an Integrating Boundary Object for Interdisciplinary Research.** *Risk Analysis* **0**.
- 49.* Verburg PH, Dearing JA, Dyke JG, Leeuw Svd, Seitzinger S, Steffen W, Syvitski J: **Methods and approaches to modelling the Anthropocene.** *Global Environmental Change* 2016, **39**:328-340.
Paper providing a broad overview of challenges in human-environmental systems modelling
50. Prestele R, Arneth A, Bondeau A, de Noblet-Ducoudré N, Pugh TAM, Sitch S, Stehfest E, Verburg PH: **Current challenges of implementing anthropogenic land-use and land-cover change in models contributing to climate change assessments.** *Earth Syst. Dynam.* 2017, **8**:369-386.
51. Song X-P, Hansen MC, Stehman SV, Potapov PV, Tyukavina A, Vermote EF, Townshend JR: **Global land change from 1982 to 2016.** *Nature* 2018, **560**:639-643.
52. Hurtt G, Chini L, Frolking S, Betts R, Feddema J, Fischer G, Fisk J, Hibbard K, Houghton R, Janetos A, et al.: **Harmonization of land-use scenarios for the period 1500–2100: 600-years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands.** *Climatic Change* 2011, **109**:117-161.
53. Lawrence PJ, Lawrence DM, Hurtt GC: **Attributing the Carbon Cycle Impacts of CMIP5 Historical and Future Land Use and Land Cover Change in the Community Earth System Model (CESM1).** *Journal of Geophysical Research: Biogeosciences* 2018, **123**:1732-1755.
54. Popp A, Calvin K, Fujimori S, Havlik P, Humpenöder F, Stehfest E, Bodirsky BL, Dietrich JP, Doelmann JC, Gusti M, et al.: **Land-use futures in the shared socio-economic pathways.** *Global Environmental Change* 2017, **42**:331-345.
55. Meyfroidt P, Roy Chowdhury R, de Bremond A, Ellis EC, Erb KH, Filatova T, Garrett RD, Grove JM, Heinimann A, Kuemmerle T, et al.: **Middle-range theories of land system change.** *Global Environmental Change* 2018, **53**:52-67.
56. Magliocca NR, Brown DG, Ellis EC: **Cross-Site Comparison of Land-Use Decision-Making and Its Consequences across Land Systems with a Generalized Agent-Based Model.** *PLOS ONE* 2014, **9**:e86179.
57. Gomiero T: **Agriculture and degrowth: State of the art and assessment of organic and biotech-based agriculture from a degrowth perspective.** *Journal of Cleaner Production* 2018, **197**:1823-1839.

58. Müller MF, Yoon J, Gorelick SM, Avisse N, Tilmant A: **Impact of the Syrian refugee crisis on land use and transboundary freshwater resources.** *Proceedings of the National Academy of Sciences* 2016, **113**:14932.
- 58a. Fischer J, Riechers M: **A leverage points perspective on sustainability.** *Nature Climate Change* 2018, **8**:109-116.
59. Verburg PH, Crossman N, Ellis EC, Heinimann A, Hostert P, Mertz O, Nagendra H, Sikor T, Erb KH, Golubiewski N, et al.: **Land system science and sustainable development of the earth system: A global land project perspective.** *Anthropocene* 2015, **12**:29-41.
60. Seppelt R, Lautenbach S, Volk M: **Identifying trade-offs between ecosystem services, land use, and biodiversity: a plea for combining scenario analysis and optimization on different spatial scales.** *Current Opinion in Environmental Sustainability* 2013, **5**:458-463.
61. Verhagen W, van der Zanden EH, Strauch M, van Teeffelen AJA, Verburg PH: **Optimizing the allocation of agri-environment measures to navigate the trade-offs between ecosystem services, biodiversity and agricultural production.** *Environmental Science & Policy* 2018, **84**:186-196.
- 62.** Wolff S, Schrammeijer EA, Schulp CJE, Verburg PH: **Meeting global land restoration and protection targets: What would the world look like in 2050?** *Global Environmental Change* 2018, **52**:259-272.
Exemplary approach of using land systems modelling to support envisioning of global sustainability targets.
63. Mehrabi Z, Ellis EC, Ramankutty N: **The challenge of feeding the world while conserving half the planet.** *Nature Sustainability* 2018, **1**:409-412.
64. Verkerk PJ, Lindner M, Pérez-Soba M, Paterson JS, Helming J, Verburg PH, Kuemmerle T, Lotze-Campen H, Moiseyev A, Müller D, et al.: **Identifying pathways to visions of future land use in Europe.** *Regional Environmental Change* 2018, **18**:817-830.
65. Rosa IMD, Pereira HM, Ferrier S, Alkemade R, Acosta LA, Akcakaya HR, den Belder E, Fazel AM, Fujimori S, Harfoot M, et al.: **Multiscale scenarios for nature futures.** *Nature Ecology & Evolution* 2017, **1**:1416-1419.
- 66.* Cooper GS, Dearing JA: **Modelling future safe and just operating spaces in regional social-ecological systems.** *Science of The Total Environment* 2019, **651**:2105-2117.
Interesting approach of integrated modelling of sustainability pathways combining exploratory and back-casting approaches.